





Accessing TMDs via single and double-spin asymmetries at HERMES

Luciano Pappalardo

pappalardo@fe.infn.it




(for the HERMES Collaboration)

Quantum phase-space tomography of the nucleon




Longitudinal momentum structure of the nucleon

Quantum phase-space tomography of the nucleon




Longitudinal momentum structure of the nucleon



QCDSF/UKQCD, PRL 98 (07)

3D picture in coordinate space



A.B., F. Conti, M. Radici, PRD78 (08)

3D picture in momentum space

The table of TMDs

momentum

helicity

transversity

		quark		
		U	L	T
n u c l e o n	U	f_1		
	L		g_1	
	T			h_1

The table of TMDs



		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_1 - h_1^\perp -

functions in red are naive T-odd

functions in green box are chirally odd

TMDs can be studied by measuring azimuthal asymmetries in SIDIS

Distribution Functions (DF)				
		quark		
		U	L	T
n u c l e o n	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_1 - h_{1T}^\perp -




functions in red are naive T-odd

functions in green box are chirally odd

TMDs can be studied by measuring azimuthal asymmetries in SIDIS

Distribution Functions (DF)				
		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp



$$\sigma^{ep \rightarrow ehX} = \sum_q DF \otimes \sigma^{eq \rightarrow eq} \otimes FF^h$$



Fragmentation Functions (FF)				
		quark		
		U	L	T
had.	U	D_1		H_1^\perp -
		Unpol. FF		Collins FF

functions in red are naive T-odd


functions in green box are chirally odd

TMDs can be studied by measuring azimuthal asymmetries in SIDIS

Distribution Functions (DF)				
		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp	g_{1T}^\perp	h_1 - h_{1T}^\perp -



$$\sigma^{ep \rightarrow ehX} = \sum_q \text{DF} \otimes \sigma^{eq \rightarrow eq} \otimes \text{FF}'$$




The SIDIS cross section exhibits asymmetries in the azimuthal angles ϕ and ϕ_S

Fragmentation Functions (FF)				
		quark		
		U	L	T
had.	U	D_1		H_1^\perp -
		Unpol. FF		Collins FF

functions in red are naive T-odd

functions in green box are chirally odd




The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x}\right)$$

$$\left\{ \begin{aligned} & \left[F_{UU,T} + \epsilon F_{UU,L} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)} \right] \\ & + \lambda_l \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right] \\ & + S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right] \\ & + S_L \lambda_l \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right] \\ & + S_T \left[\sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \right. \\ & \quad + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & \quad + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & \quad \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \right] \\ & + S_T \lambda_l \left[\sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \right. \\ & \quad + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & \quad \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \right] \end{aligned} \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_1 - h_{1T}^\perp -



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x}\right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_l \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$


$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_l \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$

$$+ S_T \lambda_l \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \left. \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x}\right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$


$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$

$$+ S_T \lambda_L \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \left. \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_1 - h_{1T}^\perp -



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h,\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$


$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$

$$+ S_T \lambda_L \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \left. \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$


$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$

$$+ S_T \lambda_L \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \left. \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_{1T}^\perp -



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)} \end{aligned} \right]$$

$$+ \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$

$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$


$$+ S_T \lambda_L \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_{1T}^\perp -

How can we disentangle all these contributions ?

EXPERIMENT: setting the proper beam and target polarization states (U, L, T)

ANALYSIS: e.g. fitting the cross section asymmetry for opposite spin states and extracting the relevant Fourier components based on their peculiar azimuthal dependences.




The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi dP_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x}\right)$$

$$\left\{ \begin{aligned} & \left[F_{UU,T} + \epsilon F_{UU,L} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)} \right] \\ & + \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right] \\ & + S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right] \\ & + S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right] \end{aligned} \right.$$

$$\left. \begin{aligned} & + S_T \left[\sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \right. \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \right] \\ & + S_T \lambda_L \left[\sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \right. \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \right] \end{aligned} \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_{1T}^\perp -



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi dP_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$

$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$


$$+ S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$

$$+ S_T \lambda_L \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \left. \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_{1T}^\perp -

See talk from [Marco Contalbrigo](#)
Wednesday morning
(WG6: Spin Physics)



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi dP_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_l \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$

$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_l \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$


$$+ S_T \lambda_l \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \Bigg\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp

Sivers

transversity
pretzelosity

$A_{UT} SSA$: transv. pol. target



The SIDIS cross-section

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi dP_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \end{aligned} \right.$$

$$+ \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right]$$

$$+ S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right]$$

$$+ S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right]$$

$$+ S_T \left[\begin{aligned} & \sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \\ & + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \end{aligned} \right]$$

$$+ S_T \lambda_L \left[\begin{aligned} & \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \end{aligned} \right] \left. \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp

Sivers


worm-gear

transversity
pretzelosity

A_{UT} SSA: transv. pol. target

A_{LT} DSA: transv. pol. target
& long. pol. beam

Selected results from A_{UT} SSAs



The Sivers effect

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$


$$\left\{ \begin{aligned} & \left[F_{UU,T} + \epsilon F_{UU,L} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)} \right] \\ & + \lambda_L \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right] \\ & + S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right] \\ & + S_L \lambda_L \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right] \\ & + S_T \left[\sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \right. \\ & \left. + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1-\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \right] \\ & + S_T \lambda_L \left[\sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \right] \end{aligned} \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp

Sivers effect


$$\propto f_{1T}^\perp(x, p_T^2) \otimes D_1(z, k_T^2)$$




- correlation between parton transverse momentum and nucleon transverse polarization
- requires orbital angular momentum



Sivers pions amplitudes

[Airapetian *et al.*, Phys. Rev. Lett. 103 (2009) 152002]



-  significantly positive
-  clear rise with z
-  rise at low $P_{h\perp}$, plateau at high $P_{h\perp}$


 slightly positive

 consistent with zero

 **First evidence of non-zero Sivers func.**

Sivers pions amplitudes

[Airapetian et al., Phys. Rev. Lett. 103 (2009) 152002]



- ☞ significantly positive
- ☞ clear rise with z
- ☞ rise at low $P_{h\perp}$, plateau at high $P_{h\perp}$
- ☞ slightly positive
- ☞ consistent with zero
- ☑ First evidence of non-zero Sivers func.

Large positive π^+ signal is dominated by scattering off u-quarks:


$$2\langle \sin(\phi - \phi_S) \rangle_{UT}^{\pi^+} \propto - \frac{f_{1T}^{\perp,u}(x, p_T^2) \otimes_W D_1^{u \rightarrow \pi^+}(z, k_T^2)}{f_1^u(x, p_T^2) \otimes D_1^{u \rightarrow \pi^+}(z, k_T^2)} \rightarrow \text{u-quark Sivers DF} < 0$$

null signal for π^- indicates that **d-quark Sivers DF > 0** (cancellation)

confirmed by phenomenological fits (Torino group) and several theoretical predictions!

Sivers kaons amplitudes

[Airapetian *et al.*, Phys. Rev. Lett. 103 (2009) 152002]



☞ significantly positive


☞ clear rise with z

☞ rise at low $P_{h\perp}$, plateau at high $P_{h\perp}$

☞ slightly positive

Sivers kaons amplitudes


[Airapetian et al., Phys. Rev. Lett. 103 (2009) 152002]



- ☞ significantly positive
- ☞ clear rise with z
- ☞ rise at low $P_{h\perp}$, plateau at high $P_{h\perp}$

- ☞ slightly positive

π^+/K^+ production dominated by u-quarks, but:




❓ $\pi^+ \equiv |u\bar{d}\rangle$, $K^+ \equiv |u\bar{s}\rangle \rightarrow$ non trivial role of sea quarks

❓ impact of different k_T dependence of FFs in the convolution integral \otimes_W

Sivers kaons amplitudes

[Airapetian et al., Phys. Rev. Lett. 103 (2009) 152002]





- ☞ significantly positive
- ☞ clear rise with z
- ☞ rise at low $P_{h\perp}$, plateau at high $P_{h\perp}$

☞ slightly positive

- each x -bin divided into two Q^2 bins
- only in low- Q^2 region significant (90% C.L.) deviation is observed

π^+/K^+ production dominated by u -quarks, but:


☞ Higher-twist contrib. for Kaons



☞ $\pi^+ \equiv |u\bar{d}\rangle$, $K^+ \equiv |u\bar{s}\rangle \rightarrow$ non trivial role of sea quarks

☞ impact of different k_T dependence of FFs in the convolution integral \otimes_W

The Collins effect




$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi dP_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & \left[F_{UU,T} + \epsilon F_{UU,L} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)} \right] \\ & + \lambda_l \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right] \\ & + S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right] \\ & + S_L \lambda_l \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right] \\ & + S_T \left[\sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \right. \\ & \quad + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \\ & \quad + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & \quad \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \right] \\ & + S_T \lambda_l \left[\sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \right. \\ & \quad + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & \quad \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \right] \end{aligned} \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp


Collins effect

- $\propto h_1(x, p_T^2) \otimes H_1^\perp(z, k_T^2)$
- correlation between parton transverse polarization in a transversely polarized nucleon and transverse momentum of the produced hadron



Collins pions amplitudes

[Airapetian *et al.*, Phys. Lett. B 693 (2010) 11-16]



- ☞ significantly positive
- ☞ clear rise with x and z

- ☞ consistent with zero

- ☞ large and negative!




$$H_1^{\perp, \text{unfav}}(z) \approx -H_1^{\perp, \text{fav}}(z)$$

☑ Non-zero Collins effect observed


☑ Both transversity and Collins function sizeable!

Consistent with Belle measurements at e^+e^- collider machines



Collins kaons amplitudes


[Airapetian *et al.*, Phys. Lett. B 693 (2010) 11-16]



- ☞ significantly positive
- ☞ clear rise with x and z
- ☞ consistent with zero

☑ Non-zero Collins effect observed

☑ Both transversity and Collins function sizeable!



The "pretzelosity"

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi dP_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & \left[F_{UU,T} + \epsilon F_{UU,L} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)} \right] \\ & + \lambda_l \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right] \\ & + S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right] \\ & + S_L \lambda_l \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right] \\ & + S_T \left[\sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi - \phi_S)} + \epsilon F_{UT,L}^{\sin(\phi - \phi_S)} \right) \right. \\ & \left. + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi + \phi_S)} + \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi - \phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi - \phi_S)} \right] \\ & + S_T \lambda_l \left[\sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi - \phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \right. \\ & \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi - \phi_S)} \right] \end{aligned} \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_{1T}^\perp - h_{1T}^\perp -


pretzelosity

$$\propto h_{1T}^\perp(x, p_T^2) \otimes H_1^\perp(z, k_T^2)$$

- characterizes the p_T dependence of the transverse quark polarization in a transversely polarized nucleon.

- can be linked to the non-spherical shape of the nucleon resulting from substantial quark orbital angular momentum


The $\sin(3\phi-\phi_S)$ Fourier component



All amplitudes consistent with zero

...suppressed by two powers of $P_{h\perp}$
w.r.t. Collins and Sivers amplitudes

New results from A_{LT} DSAs



The worm-gear g_{1T}^\perp

$$\frac{d\sigma^h}{dx dy d\phi_S dz d\phi d\mathbf{P}_{h\perp}^2} = \frac{\alpha^2 y^2}{xyQ^2 2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right)$$

$$\left\{ \begin{aligned} & [F_{UU,T} + \epsilon F_{UU,L} \\ & + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) F_{UU}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{UU}^{\cos(2\phi)}] \\ & + \lambda_l \left[\sqrt{2\epsilon(1-\epsilon)} \sin(\phi) F_{LU}^{\sin(\phi)} \right] \\ & + S_L \left[\sqrt{2\epsilon(1+\epsilon)} \sin(\phi) F_{UL}^{\sin(\phi)} + \epsilon \sin(2\phi) F_{UL}^{\sin(2\phi)} \right] \\ & + S_L \lambda_l \left[\sqrt{1-\epsilon^2} F_{LL} + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi) F_{LL}^{\cos(\phi)} \right] \\ & + S_T \left[\sin(\phi - \phi_S) \left(F_{UT,T}^{\sin(\phi-\phi_S)} + \epsilon F_{UT,L}^{\sin(\phi-\phi_S)} \right) \right. \\ & \quad + \epsilon \sin(\phi + \phi_S) F_{UT}^{\sin(\phi+\phi_S)} + \epsilon \sin(3\phi - \phi_S) F_{UT}^{\sin(3\phi-\phi_S)} \\ & \quad + \sqrt{2\epsilon(1+\epsilon)} \sin(\phi_S) F_{UT}^{\sin(\phi_S)} \\ & \quad \left. + \sqrt{2\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) F_{UT}^{\sin(2\phi-\phi_S)} \right] \\ & + S_T \lambda_l \left[\sqrt{1-\epsilon^2} \cos(\phi - \phi_S) F_{LT}^{\cos(\phi-\phi_S)} \right. \\ & \quad + \sqrt{2\epsilon(1-\epsilon)} \cos(\phi_S) F_{LT}^{\cos(\phi_S)} \\ & \quad \left. + \sqrt{2\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) F_{LT}^{\cos(2\phi-\phi_S)} \right] \end{aligned} \right\}$$

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T}^\perp -	h_1^\perp - h_{1T}^\perp -

Worm-gear




$$\propto g_{1T}^\perp(x, p_T^2) \otimes D_1(z, k_T^2)$$

- describes the probability to find longitudinally polarized quarks in a transversely polarized nucleon (\rightarrow “**trans-helicity**”)

- accessible in LT DSAs through the leading-twist $\cos(\phi - \phi_S)$ Fourier component

The worm-gear g_{1T}^\perp

- The only TMD that is both **chiral-even** and **naïve-T-even**
- requires interference between wave funct. components that differ by 1 unit of OAM



S. Boffi et al. (2009)
 Phys. Rev. D 79 094012
 Light cone constituent quark model
 flavorless
 dashed line: interf. L=0, L=1
 dotted line: interf L=1, L=2

		quark		
		U	L	T
nucleon	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_{1T}^\perp


⇒ related to quark orbital motion inside nucleons

- Many models support simple relations among g_{1T}^\perp and other TMDs:


- $g_{1T}^q = -h_{1L}^{\perp q}$ (also supported by Lattice QCD and first data)

- $g_{1T}^{q(1)}(x) \stackrel{WW-type}{\approx} x \int_x^1 \frac{dy}{y} g_1^q(y)$ (Wandzura-Wilczek appr.)

The $\cos(\phi-\phi_S)$ Fourier component New results!




The $\cos(\phi-\phi_S)$ Fourier component New results!





positive!

Resembles Hall-A results on trans. pol. ^3He targ



The $\cos(\phi-\phi_S)$ Fourier component New results!



slightly positive ?



≈ 0



positive!




slightly positive ?




≈ 0

The $\cos(\phi_S)$ Fourier component New results!



The $\cos(2\phi-\phi_S)$ Fourier component New results!






Conclusions

The existence of an intrinsic **quark transverse motion** gives origin to azimuthal asymmetries in the hadron production direction in SIDIS


- **significant Collins amplitudes observed for charged pions and K^+**
→ preliminary results enabled first extraction of transversity and Collins FF (by Torino group)
- **significant Sivers amplitudes observed for π^+ and K^+**
→ clear evidence of non-zero T-odd Sivers function
→ (indirect) evidence for non-zero quark orbital angular momentum
→ hint of non-trivial role of sea quarks and of higher-twist contrib. for positive kaons
- **first results on A_{LT} SSAs sensitive to worm-gear g_{1T}^\perp**
→ non-zero amplitudes observed for the $\cos(\phi - \phi_S)$ Fourier component for π^- (π^+, K^+)

Back-up slides


The HERMES experiment at HERA



TRD, Calorimeter, preshower, RICH:
lepton-hadron > 98%



hadron separation



$\pi \sim 98\%$, $K \sim 88\%$, $P \sim 85\%$

Accessing the polarized cross section through SSAs

Full HERMES transverse data (02-05 data with $\langle P_T \rangle \approx 73\%$)

The relevant Fourier components were extracted through a ML fit of the hadron yields for opposite target transverse spin states, alternately binned in x , z , and $P_{h\perp}$, but unbinned in ϕ and ϕ_S (\rightarrow acceptance effects on azimuthal angles cancel out)

$Q^2 > 1 \text{ GeV}^2$
$W^2 > 10 \text{ GeV}^2$
$0.023 < x < 0.4$
$y < 0.95$
$0.2 < z < 0.7$
$2 \text{ GeV} < P_h < 15 \text{ GeV}$

$$L = \prod_i^{N^h} P_i(\phi_i, \phi_{S,i}, P_{T,i}; 2\langle \sin(m\phi \pm n\phi_S) \rangle_{UT}^h) = \prod_i^{N^h} \left[1 + P_{T,i} \left(2\langle \sin(m\phi \pm n\phi_S) \rangle_{UT}^h \sin(m\phi_i \pm n\phi_{S,i}) \right) \right]$$

probability of i_{th} SIDIS event

free parameter

This is equivalent to perform a Fourier decomposition of the cross section asymmetry in the limit of vanishingly small ϕ and ϕ_S bins

$$A_{UT}^h(\phi, \phi_S) = \frac{1}{|P_T|} \frac{d\sigma^h(\phi, \phi_S) - d\sigma^h(\phi, \phi_S + \pi)}{d\sigma^h(\phi, \phi_S) + d\sigma^h(\phi, \phi_S + \pi)}$$


$$\sim \sin(\phi + \phi_S) \sum_q e_q^2 \mathcal{I} \left[\frac{k_T \hat{P}_{h\perp}}{M_h} h_1^q(x, p_T^2) H_1^{\perp,q}(z, k_T^2) \right]$$

$$+ \sin(\phi - \phi_S) \sum_q e_q^2 \mathcal{I} \left[\frac{p_T \hat{P}_{h\perp}}{M} f_{1T}^{\perp,q}(x, p_T^2) D_1^q(z, k_T^2) \right] + \dots$$



\mathcal{I} [...]: convolution integral over initial (p_T) and final (k_T) quark transverse momenta

Sivers amplitudes

[A. Airapetian et al., Phys. Rev.Lett. 94 (2005) 012002]





- First observation of T-odd Sivers effects in SIDIS!
- Main features confirmed by new high-statistics results




Collins amplitudes

[A. Airapetian et al., Phys. Rev.Lett. 94 (2005) 012002]





- ☑ First observation of non-zero Collins amplitudes
- ☑ π^+/π^- amplit. of opposite sign
- ☑ Main features confirmed by new high-statistics results



The pion-difference asymmetry



Contribution by decay of exclusively produced vector mesons (ρ^0, ω, ϕ) is not negligible (6-7% for pions and 2-3% for kaons), though substantially limited by the requirement $z < 0.7$.



a new observable

$$A_{UT}^{\pi^+ - \pi^-}(\phi, \phi_S) \equiv \frac{1}{P_T} \frac{(\sigma_{U\uparrow}^{\pi^+} - \sigma_{U\uparrow}^{\pi^-}) - (\sigma_{U\downarrow}^{\pi^+} - \sigma_{U\downarrow}^{\pi^-})}{(\sigma_{U\uparrow}^{\pi^+} - \sigma_{U\uparrow}^{\pi^-}) + (\sigma_{U\downarrow}^{\pi^+} - \sigma_{U\downarrow}^{\pi^-})}$$



Contribution from exclusive ρ^0 largely cancels out!



- significantly positive Sivers and Collins amplitudes are obtained
- measured amplitudes are not generated by exclusive VM contribution

The pion-difference asymmetry


Contribution by decay of exclusively produced vector mesons (ρ^0, ω, ϕ) is not negligible (6-7% for pions and 2-3% for kaons), though substantially limited by the requirement $z < 0.7$.



a new observable →


$$A_{UT}^{\pi^+ - \pi^-}(\phi, \phi_S) \equiv \frac{1}{P_T} \frac{(\sigma_{U\uparrow}^{\pi^+} - \sigma_{U\uparrow}^{\pi^-}) - (\sigma_{U\downarrow}^{\pi^+} - \sigma_{U\downarrow}^{\pi^-})}{(\sigma_{U\uparrow}^{\pi^+} - \sigma_{U\uparrow}^{\pi^-}) + (\sigma_{U\downarrow}^{\pi^+} - \sigma_{U\downarrow}^{\pi^-})}$$

Contribution from exclusive ρ^0 largely cancels out!



$$A_{UT}^{\pi^+ - \pi^-} = - \frac{4f_{1T}^{\perp, u_v} - f_{1T}^{\perp, d_v}}{4f_1^{u_v} - f_1^{d_v}}$$

(cancellation of FFs assuming charge-conjugation and isospin symmetry)





provides access to Sivers valence quarks distribution!

2-D Collins pions amplitudes



Kinematic dependencies often don't factorize \rightarrow correlations among variables

\rightarrow bin in as many independent variables as possible (multidim. analysis)

X vs. Z



X vs. P_{h⊥}




2-D Sivers pions amplitudes




Kinematic dependencies often don't factorize → correlations among variables

→ bin in as many independent variables as possible (multidim. analysis)

X vs. Z







X vs. $P_{h\perp}$




2-D moments for π^\pm : z vs. $P_{h\perp}$

Collins: Z vs. $P_{h\perp}$


Sivers: Z vs. $P_{h\perp}$




Collins amplitudes: twist-4 contrib ?




$\sin(\phi_S)$: Q^2 dependence for π^-



Siver amplitudes: additional studies



- ☞ No systematic shifts observed between high and low Q^2 amplitudes for both π^+ and K^+
- No indication of important contributions from exclusive VM



- ☞ test presence of $1/Q^2$ -suppressed contributions
- separate each x -bin in two Q^2 bins
- hint of higher-twist contributions to the K^+ amplitude

Probing g_{1T}^\perp through Double Spin Asymmetries

$$F_{LT}^{\cos(\phi_h - \phi_s)} = c \left[\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} g_{1T}^\perp D_1 \right]$$

$$F_{LT}^{\cos \phi_s} = \frac{2M}{Q} c \left\{ - \left(x g_T D_1 + \frac{M_h}{M} h_1 \frac{\tilde{E}}{z} \right) + \frac{\mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[\left(x e_T H_1^\perp - \frac{M_h}{M} g_{1T}^\perp \frac{\tilde{D}^\perp}{z} \right) + \left(x e_T^\perp H_1^\perp + \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{G}^\perp}{z} \right) \right] \right\}$$


$$F_{LT}^{\cos(2\phi_h - \phi_s)} = \frac{2M}{Q} c \left\{ - \frac{2(\hat{\mathbf{h}} \cdot \mathbf{p}_T)^2 - \mathbf{p}_T^2}{2M^2} \left(x g_T^\perp D_1 + \frac{M_h}{M} h_{1T}^\perp \frac{\tilde{E}}{z} \right) + \frac{2(\hat{\mathbf{h}} \cdot \mathbf{k}_T)(\hat{\mathbf{h}} \cdot \mathbf{p}_T) - \mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[\left(x e_T H_1^\perp - \frac{M_h}{M} g_{1T}^\perp \frac{\tilde{D}^\perp}{z} \right) - \left(x e_T^\perp H_1^\perp + \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{G}^\perp}{z} \right) \right] \right\}$$

Probing g_{1T}^\perp through Double Spin Asymmetries

$$F_{LT}^{\cos(\phi_h - \phi_S)} = c \left[\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} g_{1T} D_1 \right]$$

$$F_{LT}^{\cos \phi_S} = \frac{2M}{Q} c \left\{ - \left(x g_T D_1 + \frac{M_h}{M} h_1 \frac{\tilde{E}}{z} \right) + \frac{\mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[\left(x e_T H_1^\perp - \frac{M_h}{M} g_{1T} \frac{\tilde{D}^\perp}{z} \right) + \left(x e_T^\perp H_1^\perp + \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{G}^\perp}{z} \right) \right] \right\}$$


$$F_{LT}^{\cos(2\phi_h - \phi_S)} = \frac{2M}{Q} c \left\{ - \frac{2(\hat{\mathbf{h}} \cdot \mathbf{p}_T)^2 - \mathbf{p}_T^2}{2M^2} \left(x g_T^\perp D_1 + \frac{M_h}{M} h_{1T}^\perp \frac{\tilde{E}}{z} \right) + \frac{2(\hat{\mathbf{h}} \cdot \mathbf{k}_T)(\hat{\mathbf{h}} \cdot \mathbf{p}_T) - \mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[\left(x e_T H_1^\perp - \frac{M_h}{M} g_{1T} \frac{\tilde{D}^\perp}{z} \right) - \left(x e_T^\perp H_1^\perp + \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{G}^\perp}{z} \right) \right] \right\}$$



The simplest way to probe worm-gear g_{1T}^\perp is through the $\cos(\phi - \phi_S)$ Fourier component


$$2 \langle \cos(\phi - \phi_S) \rangle_{LT}^h \equiv \frac{\sqrt{1 - \varepsilon^2} F_{LT}^{\cos(\phi_h - \phi_S)}}{F_{UU,T}} = \frac{c \left[- \frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} g_{1T}^{\perp,q}(x, \mathbf{p}_T^2) D_1^q(z, z^2 \mathbf{k}_T^2) \right]}{c [f_1^q(x, \mathbf{p}_T^2) D_1^q(z, z^2 \mathbf{k}_T^2)]}$$

Model calculations




S. Boffi et al. (2009)
 Phys. Rev. D 79 094012
 Light cone constituent quark model

dashed line: interf. L=0, L=1
 dotted line: interf L=1, L=2



A. Kotzinian et al. (2006)
 arXiv:hep-ph/0603194v5

- Wandzura-Wilczek app.
- LO GRV98 (unpol. DFs)
- LO GRV2000 (pol. DFs)





H. Avakian et al. (2010)
 Phys. Rev. D 81 074035
 Bag model

B. Pasquini et al. (2008)
 Phys. Rev. D 78 034025
 Light cone constituent quark model

dashed line:
 Wandzura-Wilczek app.

R. Jakob et al.
 Nucl. Phys. A 626 937 (1997)
 Spectator model







Model predictions for $A_{LT}^{\cos(\phi-\phi_S)}$

A. Kotzinian et al. (2006)
arXiv:hep-ph/0603194v5

- Wandzura-Wilczek app.
- LO GRV98 (unpol. DFs)
- LO GRV2000 (pol. DFs)
- Kretzer FFs

Proton and deuteron targets
Similar predictions for COMPASS and JLAB




S. Boffi et al. (2009)
Phys. Rev. D 79 094012
Light cone constituent
quark model

A. Bacchetta et al. (2010)
Eur.Phys.J. A45 373-388
diquark spectator model
Pt-weighted asymmetry

J. Zhu and B0-Qiang Ma, Phys. Let. B 696 (2011) 246
Light-cone quark-diquark model

Two different prescriptions
Similar predictions for n and d targets and for
COMPASS and JLAB

Experimental status




A. Kotzinian
[arXiv:0705.2402v1 \[hep-ex\]](https://arxiv.org/abs/0705.2402v1)

COMPASS results on transversely pol. D target:

- all consistent with zero
- never published
- no results yet released on proton target


Experimental status

JLAB Neutron Transversity Collaboration
preliminary results on transversely pol. ^3He target (neutron target)



J. Huang, in *12th International Conference on Meson-Nucleon Physics and the Structure of the Nucleon* (2010), URL

		quark		
		U	L	T
n u c l e o n	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_1 h_{1T}^\perp



$$d\sigma = d\sigma_{UU}^0 + \cos 2\phi d\sigma_{UU}^1$$

$$+ \underset{L}{S} \left\{ \sin 2\phi d\sigma_{UL}^4 \right\}$$


$$+ \underset{T}{S} \left\{ \sin(\phi - \phi_s) d\sigma_{UT}^8 + \sin(\phi + \phi_s) d\sigma_{UT}^{10} \right\}$$

$$+ \frac{1}{Q}$$


$$+ \lambda_e \left[\cos(\phi - \phi_s) d\sigma_{LT}^{13} + \frac{1}{Q} \right]$$

Worm-gear (UL) (Kotzinian-Mulders)


- $\propto h_{1L}^\perp(x, p_T^2) \otimes H_1^\perp(z, k_T^2)$
- describes the probability to find transversely polarized quarks in a longitudinally polarized nucleon
- accessible in UT measurements through $\sin(2\phi + \phi_s)$ Fourier component



The $\sin(2\phi + \phi_S)$ Fourier component




- arises solely from longitudinal (w.r.t. virtual photon direction) component of the target spin



- related to $\langle \sin(2\phi) \rangle_{UL}$ Fourier comp:

$$2\langle \sin(2\phi + \phi_S) \rangle_{UT}^h \propto \frac{1}{2} \sin(\mathcal{G}_{l\gamma^*}) 2\langle \sin(2\phi) \rangle_{UL}^h$$
- sensitive to **worm-gear** h_{1L}^\perp
- suppressed by one power of $P_{h\perp}$ w.r.t. Collins and Sivers amplitudes
- **no significant non-zero signal observed (except maybe for K+)**

		quark		
		U	L	T
n u c i o n	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	h_1 h_{1T}^\perp



$$d\sigma = d\sigma_{UU}^0 + \cos 2\phi d\sigma_{UU}^1$$

$$+ \mathbf{S}_L \left\{ \sin 2\phi d\sigma_{UL}^4 + \right.$$


$$\left. + \mathbf{S}_T \left\{ \sin(\phi - \phi_s) d\sigma_{UT}^8 + \sin(\phi \right.$$

$$\left. + \frac{1}{Q} \sin(2\phi - \phi_s) d\sigma_{UT}^{11} - \frac{1}{Q} \sin \phi_s d\sigma_{UT}^{12} \right.$$


$$\left. + \lambda_e \left[\cos(\phi - \phi_s) d\sigma_{LT}^{13} + \frac{1}{Q} \right. \right.$$

Worm-gear (LT)

- $\propto g_{1T}^\perp(x, p_T^2) \otimes D_1(z, k_T^2)$
- describes the probability to find longitudinally polarized quarks in a transversely polarized nucleon
- accessible in UT measurements through sub-leading $\sin(2\phi - \phi_s)$ Fourier comp.



The subleading-twist $\sin(2\phi-\phi_S)$ Fourier component




- sensitive to worm-gear g_{1T}^\perp , Pretzelosity and Sivers function:

$$\propto W_1(p_T, k_T, P_{h\perp}) \left(x f_T^\perp D_1 - \frac{M_h}{M} h_{1T}^\perp \frac{\tilde{H}}{z} \right) - W_2(p_T, k_T, P_{h\perp}) \left[\left(x h_T H_1^\perp + \frac{M_h}{M} g_{1T} \frac{\tilde{G}^\perp}{z} \right) + \left(x h_T^\perp H_1^\perp - \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{D}^\perp}{z} \right) \right]$$

- suppressed by one power of $P_{h\perp}$ w.r.t. Collins and Sivers amplitudes


- **no significant non-zero signal observed**

The subleading-twist $\sin(\phi_S)$ Fourier component



- sensitive to worm-gear g_{1T}^\perp , Sivers function, Transversity, etc


- significant non-zero signal observed for π^- and K^- !



- low- Q^2 amplitude larger

- hint of Q^2 dependence for π^-

An alternative access to transversity: the di-hadron SSA




Di-hadron FF
(does not depend on quark transv. momentum)

Chiral-odd T-odd


Correlation between transverse spin of the fragmenting quark and the relative orbital angular momentum of the hadron pair.

Describes Spin-orbit correlation in fragmentation



azimuthal asymmetries in the direction of the outgoing hadron pairs.

azimuthal orientation of relative transv. momentum of the 2 had.



- significantly positive amplitudes
- 1st evidence of non zero dihadron FF (can be measured at e^+e^- colliders)
- independent way to access transversity
- no convolution integral involved
- limited statistical power (v.r.t. 1 hadron)